



The Challenge of Gastroparesis

A GUIDE TO UNDERSTANDING
AND MANAGEMENT



TABLE OF CONTENTS

Gastroparesis: A Serious Digestion Problem3

How the Stomach Works3

Typical Symptoms 4

Common Causes 4

When to Seek Treatment.....5

Diagnostic Testing 6

Treatment Options7

Enterra® Therapy 8

Things to Consider Before You Decide 9

Finding an Enterra Therapy Doctor 9

For More Information 10

Questions to Ask11

This guide explains what causes
gastroparesis, how to diagnose it,
and ways to treat it.

Gastroparesis: A serious digestion problem

Gastroparesis is a stomach disorder in which food moves through the stomach too slowly. In a healthy digestive system, strong muscular contractions move food through the digestive tract. With gastroparesis, the stomach muscles work poorly, or not at all, preventing the stomach from emptying properly.

Gastroparesis can be devastating. It can cause chronic nausea and vomiting and lead to malnutrition. Constant discomfort can disrupt school, work, relationships and your social life.

The number of people with gastroparesis is unknown. There is no cure, but there are therapies that improve symptoms and offer hope for a better quality of life.

How the stomach works

The stomach stores food, mixes it with stomach secretions, grinds food into small particles, and moves these particles into the small intestine. Emptying is controlled by muscle contractions in different regions of the stomach. Ideally, these processes should occur at a rate that makes digestion easy and efficient.

If stomach muscle contractions stop working properly, the result can be delayed gastric emptying. Undigested food and fluids sit in the stomach, causing chronic nausea and vomiting.



Typical symptoms

People with gastroparesis experience a number of gastrointestinal symptoms including:

Chronic nausea	Frequent vomiting	Abdominal bloating	Abdominal pain
Vomiting on an empty stomach	Dehydration	Heartburn or Gastroesophageal Reflux Disease (GERD)	Changes in blood sugar levels
Malnutrition	Electrolyte imbalances	Lack of appetite	Excessive weight loss
Feelings of fullness after only a few bites of food (early satiety)			

Nausea and vomiting have been reported as the most bothersome symptoms of gastroparesis¹

Common causes

Damage to the vagus nerve may be a cause of gastroparesis. This nerve stretches from the brain stem to the colon and controls the functions of the esophagus, stomach, and intestines. Damage to stomach muscles is another cause. There are many ways the vagus nerve or stomach muscles can be damaged:^{1,2}

DIABETES, EITHER TYPE 1 OR TYPE 2

High blood sugar can damage the vagus nerve over time. Unfortunately, gastroparesis makes blood sugar levels harder to control.

POSTSURGICAL COMPLICATIONS

Surgery on the esophagus, stomach, or upper part of the small intestine can injure the vagus nerve. Symptoms may appear immediately or years later.*

OTHER CONDITIONS AND CAUSES

Parkinson's disease, vascular disease, and pseudo-obstructions can all cause gastroparesis. Medications can also slow stomach emptying, resulting in symptoms that mimic gastroparesis.*

UNKNOWN

Frustratingly, the cause may not be known (idiopathic). Delayed gastric emptying without apparent cause or underlying abnormality is a common diagnosis.^{1,2} Rarely, gastroparesis develops after a person recovers from the flu or other viral illness.

* Enterra Therapy is not indicated for gastroparesis symptoms due to postsurgical complications, Parkinson's disease, vascular disease or pseudo-obstructions.

When to seek treatment

Many people experience occasional nausea and vomiting. Warning signs of gastroparesis include:

- Nausea and/or vomiting occurring several times a day
- Nausea and/or vomiting episodes over 12 months or more
- Weight loss over the past year
- Excessive weight gain over the past year due to improper nutrition
- Supplemental nutrition needed
- Repeated hospitalizations
- Inadequate relief of symptoms from medications
- Diminished quality of life



Diagnostic testing

Nausea and vomiting are symptoms of several other digestive disorders. After taking a complete medical history and performing a physical examination, your doctor will ask for some additional tests in order to determine a diagnosis.

PRIMARY TESTS (REQUIRED)

Upper Gastrointestinal Endoscopy

This test excludes other conditions (e.g., ulcers, physical abnormalities, or mechanical obstruction) that may be causing symptoms. A thin, flexible tube (endoscope) with a tiny camera at the tip is passed through the mouth and guided into the stomach while you are asleep. The endoscope allows the doctor to look into the stomach and at its lining.

Gastric Emptying Test (GET)

This procedure measures the speed at which food empties from the stomach into the small intestine. The GET, considered the most accurate way to diagnose gastroparesis, requires that you eat a meal in which a solid food (usually eggs or oatmeal) contains a small and safe amount of a substance that will appear on a medical scan. A scanner is placed over the abdomen to monitor the food's movement through the digestive tract.




SECONDARY TESTS (MAY NOT BE NECESSARY)

Gastroduodenal Manometry

In this test, the doctor passes a thin, pressure-sensitive tube through the mouth and into the stomach and small intestine. A computer connected to the tube monitors the strength and coordination of stomach muscle contractions and how well the stomach is emptying.

Electrogastrography (EGG)

This procedure records the electrical signals in the stomach before and after eating. An irregular electrical rhythm may indicate the presence of gastroparesis.



Treatment options

There is no cure for gastroparesis. Several treatments are available that improve symptoms and offer a more comfortable lifestyle:³

NUTRITION EDUCATION AND DIET MODIFICATIONS

This can help reduce symptoms and maintain adequate fluids and nutrition. A modified diet typically consists of liquids, restricted fats and plant fiber, and frequent, small meals.

PROKINETIC DRUGS can improve the rate of stomach emptying and thereby reduce nausea and vomiting.

ANTIEMETIC DRUGS can control nausea and vomiting, but have no effect on stomach emptying.³

ENTERAL NUTRITION

This therapy involves the delivery of liquid nutrients via a tube placed directly into the stomach or small intestine. Feeding tubes are usually temporary and used only when gastroparesis is severe.



TOTAL PARENTERAL NUTRITION (TPN)

TPN supplies nutrients to your bloodstream through an intravenous (IV) infusion. TPN is used only if enteral feeding is not tolerated or is insufficient to meet caloric needs.

SURGERY

Surgery is considered a last resort. One surgery, called pyloroplasty, involves widening the muscle that separates the stomach from the upper region of the small intestine or duodenum. Another surgery, called gastrectomy, is a medical procedure where all or part of the stomach is surgically removed.

ENTERRA® THERAPY*

This is a surgical option for patients with chronic, drug-refractory (resistant to medication) nausea and vomiting due to gastroparesis of diabetic or idiopathic (unknown) origin in patients aged 18 to 70 years. The Enterra Therapy System includes a battery-powered neurostimulator implanted beneath the skin. Your doctor can determine if you are a candidate for Enterra Therapy.

Any combination of diet modification, medication, nutritional support, surgery, and Enterra Therapy may be necessary to control symptoms of gastroparesis.

*Humanitarian Device. Authorized by Federal law for use in the treatment of chronic intractable (drug refractory) nausea and vomiting secondary to gastroparesis of diabetic or idiopathic etiology in patients aged 18 to 70 years. The effectiveness of this device for this use has not been demonstrated. Enterra Therapy received Humanitarian Device Exemption (HDE) approval from the U.S. Food and Drug Administration (FDA) in 2000.

Enterra Therapy

Gastric Electrical Stimulation for Gastroparesis

Enterra Therapy* is designed to improve chronic, medication resistant nausea and vomiting associated with gastroparesis of diabetic or idiopathic (unknown) origin.

The Enterra Therapy System consists of a neurostimulator (about the size of a pocket watch) implanted beneath the skin, usually in the lower abdominal region. Two leads (insulated wires) are implanted in the muscle wall of the stomach and connected to the neurostimulator. The procedure is performed under general anesthesia.

The neurostimulator sends mild pulses through the leads to stimulate the nerves and smooth muscles of the lower stomach. This may help control chronic nausea and vomiting.

During an office visit, the doctor adjusts the neurostimulator to customize therapy for you. The therapy can be turned off by the doctor at any time.



Over 15,000 people
have received Enterra
Therapy to help resume
everyday activities, like
taking their seat back
at the table.

*Humanitarian Device. Authorized by Federal law for use in the treatment of chronic intractable (drug refractory) nausea and vomiting secondary to gastroparesis of diabetic or idiopathic etiology in patients aged 18 to 70 years. The effectiveness of this device for this use has not been demonstrated. Enterra Therapy received Humanitarian Device Exemption (HDE) approval from the U.S. Food and Drug Administration (FDA) in 2000.

A small medical device called a neurostimulator is implanted under the skin, usually in the lower abdominal region. Two insulated wires called leads are implanted in the stomach wall muscle and then connected to the neurostimulator.



The device that appears in this graphic is a representation of the device. The graphic does not include certain identifying information that may appear on actual devices such as model number or serial number.

Things to consider before you decide

If your doctor determines that you are a candidate, please consider the following:

- The therapy is intended to reduce symptoms of chronic nausea and vomiting associated with gastroparesis. But, it is not a cure.
- Improvements in symptoms may be gradual or immediate. Results vary from person to person.
- Implanting an Enterra Therapy System has risks and side effects. Surgical complications are possible and may include infection, bleeding, bruising, and pain at the implant site. Once implanted, the system may become infected, devices may move or wear through the skin, the lead may entangle with or obstruct the bowel, lead may perforate your stomach, or irritation/inflammation over the implant site may occur. The system could stop suddenly because of mechanical or electrical problems. Any of these situations may require additional surgery or cause return of symptoms, and some can be life threatening.
- Most often, a combination of Enterra Therapy, diet modification, and medication is necessary to effectively control symptoms of gastroparesis.
- If you have an Enterra Therapy System implanted, some precautions are necessary around certain electrical and medical equipment and when going through theft detection and security screening gates.
- Enterra Therapy is not appropriate for patients who are not candidates for surgical procedures and/or anesthesia because of physical or mental conditions.
- Enterra Therapy should not be used for patients who will be exposed to diathermy (deep heat treatment).
- Enterra Therapy has not been evaluated in pregnant women, or in patients younger than age 18 or older than age 70.



Finding an Enterra Therapy provider

Visit enterramedical.com/find-a-provider to find a provider who specializes in Enterra Therapy.

Next Steps

1. Get evaluated

Your doctor will determine the appropriate screening tools and diagnostic tests for you.

2. Review the results

Review the results from screening and testing with your doctor.

3. Discuss treatment options

Ask your doctor for a treatment recommendation. Ask questions and review possible side effects of each therapy. Share any concerns you may have with your doctor.

4. Take action

With your doctor's help, choose the treatment option that is best for you. Follow the therapy plan, including diet modification, regular follow-up visits, and other instructions.

5. Seek Prior Authorization

If your doctor determines Enterra Therapy is right for you, please ask them to obtain prior authorization from your health insurance company.



For more information

PUBLICATIONS

The Official Patient's Sourcebook on Gastroparesis: A Revised and Updated Directory for the Internet Age.

Icon Health Publications, 2002

Gastroparesis – A Medical Dictionary, Bibliography, and Annotated Research Guide to Internet References.

Icon Health Publications, 2004

WEBSITES FOR EDUCATION, INFORMATION AND SUPPORT

**American College of
Gastroenterology**

www.gi.org/topics/gastroparesis/

**Association of Gastrointestinal
Motility Disorders, Inc.**

www.agmdhope.org

**Cyclic Vomiting Syndrome
Association**

www.cvsaonline.org

**Gastroparesis Patient Association
for Cures & Treatments, Inc.**

www.g-pact.org

**International Foundation for
Functional Gastrointestinal Disorders**

www.iffgd.org

**National Institute of Diabetes &
Digestive & Kidney Diseases**

[www.niddk.nih.gov/health-information/
digestive-diseases/gastroparesis](http://www.niddk.nih.gov/health-information/digestive-diseases/gastroparesis)

National Pancreas Foundation

www.pancreasfoundation.org

Oley Foundation (Nutritional Support)

www.oley.org



Visit www.enterramedical.com/introduction-to-enterra-therapy to learn more about gastroparesis and Enterra Therapy by watching a 3-minute video.

QUESTIONS TO ASK

I have some warning signs that indicate gastroparesis.

What testing do you recommend for further evaluation?

What medical conditions or medications cause symptoms of gastroparesis?

What are the potential side effects of each treatment option for gastroparesis?

What are the potential risks, complications, and side effects of the Enterra Therapy System?

What additional information can you give me about Enterra Therapy?

I have other questions and concerns (please list):


This image shows a single sheet of white paper with horizontal ruling lines. The lines are evenly spaced and run across the width of the page. There are no margins, text, or other markings on the paper.

A high-angle, top-down photograph of a woman with short brown hair, wearing a vibrant green short-sleeved button-down shirt. She is seated at a dark wooden table outdoors, with her eyes closed and a gentle smile on her face. The table is set with a meal including a large plate of food with potatoes and vegetables, a smaller plate of salad, a pitcher of water, several glasses, and a bottle of olive oil. To her right, another person wearing a light-colored shirt and a straw hat is partially visible. The background consists of dry, golden-brown grass, suggesting a rural or outdoor setting. A large, semi-circular yellow graphic element is positioned in the bottom left corner, containing white text.

To learn more about
Enterra Therapy visit
www.enterramedical.com



Enterra Medical is dedicated to helping more people with chronic gastroparesis live better lives through advancing technology, bolstering clinical science, and accelerating patient access to Enterra Therapy.



The information provided in this brochure is for general educational purposes only and is not a substitute for professional medical advice, diagnosis or treatment. Always talk to your doctor about the best treatment options for your individual situation.

1. Soykan I, Sivri B, Sarosiek I, et al. Demography, clinical characteristics, psychological and abuse profile, treatment and long-term follow-up of patients with gastroparesis. *Dig Dis Sci*. 1998;43:2398-2404.
2. Ye Y, Yin Y, Huh SY, Almansa C, Bennett D, Camilleri M. Epidemiology, Etiology, and Treatment of Gastroparesis: Real-World Evidence From a Large US National Claims Database. *Gastroenterology*. 2022;162(1):109-121.e5. doi:10.1053/j.gastro.2021.09.064.
3. American Gastroenterology Association. Technical review on the diagnosis and treatment of gastroparesis. *Gastroenterology*. 2004;127:1592-1622.

Important Safety Information

Intended Use: The Enterra® Therapy System is an implanted device that provides gastric stimulation to treat chronic, intractable, nausea and vomiting that is not well treated by drugs or other means in patients aged 18 to 70 years caused by diabetes or an unknown origin.

Contraindications: Enterra Therapy is only for patients who are healthy enough for surgical procedures and/or anesthesia. Once implanted, patients need to avoid diathermy, which is deep heat treatment from electromagnetic energy, as it may cause injury or device failure.

Warnings: Enterra Therapy has not been studied in pregnant women, patients under the age of 18, or over 70. Issues may occur if the system interacts with other implanted devices such as pacemakers. Patient injury or device failure may be caused by other medical treatments such as electrocautery, defibrillation/cardioversion, therapeutic ultrasound, or radiofrequency (RF)/microwave ablation. Patient activities may cause shocking or jolting sensations.



The Enterra II System is MR Conditional. This means that patients with the Enterra II System can safely have MRI examinations of some body parts under certain conditions. Consult your doctor to determine if you are eligible for MRI examination.

Risks: Potential risks include infection, pain at the surgery site, allergic or immune system response, lead and bowel twist together, device wearing through the skin, bruising, bleeding, loss of therapeutic effect, jolting, shocking, burning sensation, gastrointestinal or stomach issues, loss of therapy due to component failure or battery wear out, or perforated stomach which may cause life-threatening blockage or infections that require immediate medical attention including surgery. Risks can be minimized by avoiding activities such as sudden, excessive, or repetitive bending, twisting, bouncing, or stretching.

Humanitarian Device: Authorized by Federal law for the intended use described above. The effectiveness of this device has not been demonstrated.

Always discuss potential risks and benefits of the device with your physician.

For further information, please contact Enterra Medical at info@enterramedical.com.

Rx Only.

www.enterramedical.com

Enterra® is a registered trademark of Enterra Medical, Inc. in the US, EU, and other regions.

©2025 Enterra Medical, Inc. All rights reserved.

MKT-B-03254, Rev A

